



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Cost of dispersal in a social mammal: body mass loss and increased stress

Maag, Nino ; Cozzi, Gabriele ; Bateman, Andrew ; Heistermann, Michael ; Ganswindt, André ; Manser, Marta ; Clutton-Brock, Tim ; Ozgul, Arpat

Abstract: Dispersal is a key process influencing the dynamics of socially and spatially structured populations. Dispersal success is determined by the state of individuals at emigration and the costs incurred after emigration. However, quantification of such costs is often difficult, due to logistical constraints of following wide-ranging individuals. We investigated the effects of dispersal on individual body mass and stress hormone levels in a cooperative breeder, the meerkat (*Suricata suricatta*). We measured body mass and faecal glucocorticoid metabolite (fGCM) concentrations from 95 dispersing females in 65 coalitions through the entire dispersal process. Females that successfully settled lost body mass, while females that did not settle but returned to their natal group after a short period of time did not. Furthermore, dispersing females had higher fGCM levels than resident females, and this was especially pronounced during the later stages of dispersal. By adding information on the transient stage of dispersal and by comparing dispersers that successfully settled to dispersers that returned to their natal group, we expand on previous studies focusing on the earlier stages of dispersal. We propose that body mass and stress hormone levels are good indicators to investigate dispersal costs, as these traits often play an important role in mediating the effects of the environment on other life-history events and individual fitness.

DOI: <https://doi.org/10.1098/rspb.2019.0033>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-168418>

Journal Article

Accepted Version

Originally published at:

Maag, Nino; Cozzi, Gabriele; Bateman, Andrew; Heistermann, Michael; Ganswindt, André; Manser, Marta; Clutton-Brock, Tim; Ozgul, Arpat (2019). Cost of dispersal in a social mammal: body mass loss and increased stress. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 286(1896):20190033.

DOI: <https://doi.org/10.1098/rspb.2019.0033>

PROCEEDINGS OF THE ROYAL SOCIETY B

BIOLOGICAL SCIENCES

Cost of dispersal in a social mammal – body mass loss and increased stress

Journal:	<i>Proceedings B</i>
Manuscript ID	RSPB-2019-0033.R1
Article Type:	Research
Date Submitted by the Author:	16-Jan-2019
Complete List of Authors:	Maag, Nino; University of Zurich, Department of Evolutionary Biology and Environmental Studies; Cozzi, Gabriele; Zurich University, Institute of Evolutionary Biology and Environmental Studies Bateman, Andrew; University of Alberta, Biology/Mathematical and Statistical Sciences; University of Toronto, Ecology and Evolutionary Biology Heistermann, Michael; German Primate Center, Endocrinology Laboratory Ganswindt, Andre; University of Pretoria, Anatomy and Physiology Manser, Marta; University of Zurich, Institute of Evolutionary Biology and Environmental Studies Clutton-Brock, Tim; University of Cambridge, Zoology Ozgul, Arpat; University of Zurich, Institute of Evolutionary Biology and Environmental Studies
Subject:	Ecology < BIOLOGY
Keywords:	Dispersal stage, Glucocorticoid metabolites, Individual trait, Life history, Meerkat
Proceedings B category:	Ecology

SCHOLARONE™
Manuscripts

Cost of dispersal in a social mammal – body mass loss and increased stress

Nino Maag^{1,2}, Gabriele Cozzi^{1,2}, Andrew Bateman³, Michael Heistermann⁴, André Ganswindt⁵, Marta Manser^{1,2}, Tim Clutton-Brock^{2,5,6}, Arpat Ozgul^{1,2}

¹ Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

² Kalahari Research Centre, Kuruman River Reserve, Van Zylsrus 8467, South Africa

³ Department of Geography, University of Victoria, 3800 Finnerty Road, Victoria V8P 5C2, Canada

⁴ Endocrinology Laboratory, German Primate Center, Leibniz Institute for Primate Research, Kellnerweg 4, D-37077 Göttingen, Germany

⁵ Mammal Research Institute, University of Pretoria, cnr Lynnwood Road and Roper Street, Hatfield 0028, South Africa

⁶ Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, United Kingdom

Corresponding author: Nino Maag, nino.maag@gmail.com

Author contributions: NM, GC, and AO conceived the study. NM collected data on dispersing individuals, performed statistical modelling, and wrote the manuscript. AB and GC consulted on statistical modelling. AG and MH led steroid extraction and analysis. TCB and MM led the collection of data on resident groups. All authors substantially contributed to the final draft.

Data accessibility statement: Data available from the Dryad Digital Repository <https://datadryad.org/review?doi=doi:10.5061/dryad.g44j6g2>

25 **Abstract**

26 Dispersal is a key process influencing the dynamics of socially and spatially
27 structured populations. Dispersal success is determined by the state of individuals at
28 emigration and the costs incurred after emigration. However, quantification of such
29 costs is often difficult, due to logistical constraints of following wide-ranging
30 individuals. We investigated the effects of dispersal on individual body mass and
31 stress hormone levels in a cooperative breeder, the meerkat (*Suricata suricatta*). We
32 measured body mass and faecal glucocorticoid metabolite (fGCM) concentrations
33 from 95 dispersing females in 65 coalitions through the entire dispersal process.
34 Females that successfully settled lost body mass, while females that did not settle but
35 returned to their natal group after a short period of time did not. Furthermore,
36 dispersing females had higher fGCM levels than resident females, and this was
37 especially pronounced during the later stages of dispersal. By adding information on
38 the transient stage of dispersal and by comparing dispersers that successfully settled
39 to dispersers that returned to their natal group, we expand on previous studies
40 focusing on the earlier stages of dispersal. We propose that body mass and stress
41 hormone levels are good indicators to investigate dispersal costs, as these traits often
42 play an important role in mediating the effects of the environment on other life-
43 history events and individual fitness.

44 **Keywords:** Dispersal stage, glucocorticoid metabolites, individual trait, life history,
45 meerkat

46 **Introduction**

47 Dispersal is an important life-history process typically consisting of three stages:
48 emigration from a natal territory, transience through an unfamiliar landscape, and
49 settlement in a new area [1,2]. At each stage, dispersing individuals incur costs
50 resulting from different individual, social, and environmental factors, and these costs
51 influence the outcome of dispersal and consequently the associated population
52 dynamics [1,3]. Costs of dispersal have been typically measured in the form of
53 changes in survival and reproductive rates [3–5]. However, the effects of social and
54 environmental factors on survival and reproduction are often mediated by
55 morphological and physiological traits [6–8]. As such, a more mechanistic
56 understanding of dispersal costs can be achieved by investigating traits such as body
57 mass and stress hormone levels, which are known to correlate with individual fitness
58 [9,10].

59 A variety of processes can be expected to influence body mass and stress hormone
60 levels during dispersal [3]. For instance, unfamiliarity with a new area may result in
61 reduced feeding efficiency [11], which may lead to loss in body mass and increased
62 stress hormone levels. Unfamiliar territory often comes with higher susceptibility to
63 predators, which might be reflected in further elevated stress levels [12]. Furthermore,
64 dispersers are often exposed to aggression from resident conspecifics [13,14], and
65 attacks may lead to wounds and increased stress [15–17]. An overall deterioration in
66 body condition can in turn lead to a decrease in immune defence [18–20]. Social
67 factors associated with dispersal, such as loss of social rank and loss of benefits
68 rendered by group membership (e.g., secured foraging territory and anti-predator
69 vigilance), may further exacerbate morphological and physiological costs of dispersal
70 [21,22].

71 Due to the difficulties of following wide-ranging individuals in the wild,
72 quantification of morphological and physiological costs of dispersal has proven
73 challenging, and there have been only a few such studies on vertebrates. For example,
74 in the red squirrel (*Tamiasciurus hudsonicus*), where juveniles make temporary forays
75 outside the natal territory prior to emigration, no strong relationship between
76 maximum foray distance and body mass was observed [23]. In the cooperatively
77 breeding pied babbler (*Turdoides bicolor*), lone evicted individuals (“floaters”

without territory) lost more body mass than residents due to increased time allocated to vigilance and less efficient foraging [24]. In the cooperatively breeding meerkat (*Suricata suricatta*), where subordinate males undertake extraterritorial prospecting trips [25] and subordinate females are evicted from their natal group [26], both sexes experienced increased stress levels and body mass loss while outside their natal group [16,25,27]. In addition, female meerkats showed reproductive down-regulation [16] and males showed increased testosterone levels [27]. All the above-mentioned studies followed individuals during exploratory movements and early phases of dispersal. Such results are, however, not generalizable among dispersal stages, because disperser candidates are often young and inexperienced and exploratory movements are typically carried out close to the natal range [16,28].

Social species may be able to alleviate some of the costs of dispersal by forming multiple-member dispersing coalitions [29–31], and in many cooperative breeders, subordinate helpers disperse in multiple-member coalitions [13,32–34]. Larger coalitions are likely to experience reduced individual predation risk [30,35], have increased competitive ability [36–38], and faster group-size augmentation, as having more helpers increases reproductive success [31]. Dispersing in larger coalitions may also reduce deterioration in body condition. For example, meerkats that left their natal group in larger coalitions had higher hourly foraging mass gain and lower parasite loads and stress levels [38]. However, how body condition changes with varying coalition size during the entire dispersal event has not yet been formally tested.

Meerkats live in groups of 2–50 individuals and groups are characterized by the presence of a dominant pair that monopolises reproduction [26,39]. During her pregnancy, the dominant female often evicts one or multiple subordinate females [16,26]. When several females are evicted simultaneously, they usually form same-sex multiple-member dispersing coalitions [26]. After eviction, females remain within the territory of the natal group for a variable period (hereafter referred to as the “post-eviction stage”). At the end of this period, individuals are either accepted back to the natal group as “returners”, or permanently emigrate and enter transience as “emigrants”. Emigration is here defined as the time when emigrants leave the natal territory and enter transience. Returners do not leave the natal territory nor enter the transient stage [40]. Evicted females’ decision to return or emigrate is dependent on a

110 multitude of factors such as natal group size, age, rainfall, population density,
111 coalition size, and availability of unrelated males from other groups [40,41].

112 In this study, we explore the morphological and physiological costs of dispersal
113 throughout the entire dispersal process from the eviction of subordinate female
114 meerkats until either establishment of a new group or return to the natal group. As
115 such, we extend previous studies by comparing dispersers that successfully settled to
116 dispersers that returned, and by formally assessing costs during the transient stage.
117 We specifically investigate (1) how the loss in body mass and faecal glucocorticoid
118 metabolite levels vary among emigrants, returners, and resident subordinates; and (2)
119 how these measures change between emigration and transient stages. We further
120 investigate (3) how these changes vary with the size of the dispersing coalition while
121 controlling for additional individual and environmental factors.

122 **Methods**

123 We conducted our study between September 2013 and March 2017 at the Kalahari
124 Meerkat Project (KMP) located on the Kuruman River Reserve (26° 59' S, 21° 50'
125 E), South Africa. The region is characterized by low seasonal rainfall between
126 October and April and large daily and seasonal temperature variations [26].
127 Temperature data were collected hourly and precipitation data daily with an on-site
128 weather station, allowing for accurate investigation of climatic conditions.

129 *GPS data collection and definition of dispersal stages*

130 We fitted lightweight GPS radio-collars (<25 g, ~3.5% of body mass) on subordinate
131 females a few days prior to, or immediately after, eviction from their natal groups.
132 The collars were composed of a VHF module (Holohil Systems Ltd., Canada) and a
133 GPS module (CDD Ltd., Greece). Collars of this size and weight do not affect
134 meerkat behaviour and survival [42], and we did not observe any sign of distress in
135 animals carrying collars. Typically, only one individual in each dispersing coalition
136 was fitted with a radio-collar. To mount the collars, we sedated individuals using a
137 mixture of isoflurane and oxygen in compliance with the KMP protocol and in
138 collaboration with trained project staff [42]. All necessary permits to handle and tag
139 meerkats were granted to the KMP by the Department of Environment and Nature
140 Conservation of the Northern Cape, South Africa, and the Animal Ethics Committee

of the University of Pretoria (permit 'FAUNA 192/2014'). We set the collars to automatically record and store GPS locations at hourly intervals during daytime [43]. However, due to miniaturization of the GPS antenna, we observed a considerable amount of missing GPS fixes (i.e., on average 40 % of fixes were not recorded).

We used the net squared displacement (NSD) approach, which is stable against missing GPS fixes, to identify time of emigration and time of settlement. The NSD measures the square of the Euclidean distance from the place of eviction to any given GPS location along the dispersal path [44]. By visually investigating the NSD plots of each dispersal event, we identified the inflection points [45], which corresponded to time of emigration and time of settlement (Supplementary material: Figure S1). We further used field observations to validate the NSD approach.

Field observations

We located collared dispersing females by means of VHF radio-tracking every two to seven days. Study animals were part of the long-term research at the KMP and were habituated to the presence of humans [46]. At each visit, we recorded dispersing coalition size, number of associated unrelated males from other groups, and pregnancy status, and we measured individual body mass and collected faecal samples for stress hormone metabolite analysis (see below for more details). To compare dispersers (emigrants and returners) to resident females, we included data collected on same-age subordinate females that resided in the dispersers' respective natal groups (hereafter referred to as "residents").

Resident groups were located by means of VHF radio-tracking (one individual per group was carrying a radio-collar) several times each week by volunteers working at the KMP, and data on group size and composition, female pregnancy status, body mass, and faecal samples were collected. Information on birth date and relatedness of each individual (i.e., dispersers and resident) were available from the long-term database. We considered data on resident individuals for the exact period during which dispersers were absent from the group.

Determination of body mass

170 Individuals in our study population were trained to stand on an electronic balance
171 [46]. We weighted dispersing females at each visit, either in the morning before
172 foraging started or in the evening after foraging ended. To match mass measures
173 collected in the evening and morning, we subtracted the average overnight mass loss
174 (28 g for dispersers) from evening masses and treated them as morning masses on the
175 following day. We calculated average overnight mass loss by subtracting evening and
176 morning masses of dispersers for which consecutive measures were available.
177 Morning measures for resident females were more abundant as resident groups were
178 visited five times per week, and therefore, morning measures alone were sufficient for
179 residents. We excluded from the analyses mass data from pregnant females between
180 the day of conception and the day of parturition or abortion. We determined
181 pregnancy onset by backdating 28 days from the first evidence of abdominal swelling
182 [47,48]. We identified parturition and abortion from a sudden change in abdominal
183 shape and mass loss. After filtering data, we had 192 mass measurements from 25
184 emigrants, 200 from 28 returners, and 504 from 49 resident subordinates.

185 *Faecal glucocorticoid metabolite (fGCM) analysis*

186 We opportunistically collected faecal samples for fGCM analysis immediately after
187 defecation and assigned them to identified individuals. The average time lag between
188 a perceived stressor and its reflection in altered meerkat fGCM concentrations is
189 about 24 hours [49] and we therefore did not collect faecal samples within 2 days
190 after collar deployment. We kept collected samples in insulated thermal bags filled
191 with ice packs until the samples were frozen at -80°C within three hours from
192 collection. Overall, we collected 125 samples from 32 emigrants, 113 samples from
193 25 returners, and 111 samples from 47 resident subordinates. For steroid extraction,
194 we lyophilized and pulverized faecal samples, and subsequently extracted 0.10-0.11 g
195 of faecal powder with 3 ml of 80 % watery methanol [50]. We subsequently analysed
196 faecal extracts for immunoreactive glucocorticoid metabolite concentrations using a
197 group-specific enzyme immunoassay (EIA) for the measurement of 11 β -
198 hydroxyetiocholanolone [51]. The assay has previously been validated for assessing
199 glucocorticoid output in captive meerkats [49] and has recently been applied
200 successfully to study adrenocortical activity of wild meerkats [52]. Details on assay
201 performance and characteristics, including full descriptions of the assay components

and cross-reactivities are provided elsewhere [51,53]. The sensitivity of the EIA at 90 % binding was 0.6 pg. Intra- and inter-assay coefficients of variation of high- and low-value quality controls were 6.2 % (high) and 8.1 % (low) and 7.3 % (high) and 8.7 % (low), respectively. All fGCM concentrations are expressed as hormone mass per dry faecal mass.

Statistical modelling

We used linear mixed effects models to analyse variation in (1a) body mass loss and (1b) fGCM concentration among emigrants, returners, and residents. We used two additional models to compare (2a) body mass loss and (2b) fGCM concentration between post-eviction (time between eviction and emigration) and transient (time between emigration and settlement) stages. For models 2a and 2b we only used data collected on emigrants. We conducted all statistical analyses using the library *lme4* [54] in the software R [55]. We used the library *MuMin* [56] to test different combinations of individual, social, and environmental variables using model selection based on Akaike's Information Criterion [57]. As we could not identify any single best model, we calculated the natural model average across all models with ΔAIC_c values within 2 units [58]. Details for full models, descriptions of model variables, and outcomes of model selection are given in the supplements (Tables S1-S4). To ensure that predictor variables were not correlated with each other, we calculated variance inflation factors [59] for coefficients in the full models. We standardised continuous variables across all data points used for a given model by subtracting their mean and then dividing by their standard deviation. We created residuals plots (Figure S2) and investigated them visually to ensure that model assumptions were met; we did not find any considerable departure from theoretical expectations. Additionally, for each female, we obtained autocorrelation function plots from model residuals to test for temporal autocorrelation [60]. Model residuals showed minor autocorrelation in 3.8 % of the cases and we therefore did not consider temporal autocorrelation as reason for concern.

1a) Body mass in emigrants, returners, and residents: We compared the daily proportional change in body mass of emigrants to those of returners and residents. We used a proportional rather than absolute mass change because a given mass loss (or gain) in a light individual would not be equivalent to the same mass loss (or gain) in a

234 heavy individual. Because mass data were collected opportunistically, time between
 235 mass measures varied considerably and ranged from 1–81 days (average: 3.21 days,
 236 interquartile range: 1–4 days). As we could not assume linearity in mass change
 237 across time, we could not reliably and consistently calculate a typically used daily
 238 growth rate. Instead, we used a net proportional daily growth rate since eviction ($(m_t -$
 239 $m_0)/(m_0 * \Delta t)$, where m_0 is the mass at eviction and Δt is the number of days elapsed
 240 between eviction and a given mass measure m_t . Because we were interested in
 241 comparing emigrants, returners, and residents, we included a categorical variable
 242 (*strategy*) with these three levels. As larger individuals usually have higher metabolic
 243 rates and may lose proportionally more of their body mass [61], we added mass at
 244 eviction m_0 as an explanatory variable. We treated m_0 as a surrogate for body
 245 condition because we could not measure body size in the field. We also included the
 246 number of days since eviction (Δt) to investigate if time away from the group
 247 influenced mass loss. We included both *age* and $age^{1/2}$ to account for known
 248 nonlinearity in meerkat growth [62]. We included a continuous variable (*#female*)
 249 indicating the average number of females in a coalition. The reason for averaging
 250 being that only 29% of the coalitions changed in size and such changes were of only
 251 ± 1 individual in most cases. Without averaging, these coalitions would have had a
 252 disproportionate high weight in the model output. We added additional covariates,
 253 which are known to influence mass of meerkats: number of nutritionally dependent
 254 offspring younger than three months in the relevant group (*#pup*), maximum daily
 255 temperature (*temp*), and cumulative amount of rain during three months prior to mass
 256 measurements (*rain*, [62]). We assigned a unique coalition ID to each dispersal event,
 257 and resident individuals that were in dispersers' respective natal group during a given
 258 dispersal event.

259 We treated individual (*indID*) nested in coalition (*coalID*) as the random intercept
 260 terms, to account for consistency across repeated measures. Because *coalID* was
 261 specific to a given year and therefore partially captured the year effect, we did not
 262 include year as a random term to avoid overfitting. We assessed the variability within
 263 and between individuals and coalitions by calculating the within-individual and
 264 within-coalition repeatability [63]. We calculated repeatability for all models
 265 described below.

1b) *fGCM in emigrants, returners, and residents*: The distribution of *fGCM* levels was right-skewed, and we used $\log(fGCM)$ as response variable in order to assess differences in stress hormone levels between emigrants, returners, and residents. As described above, we included *strategy* as well as Δt , *age*, *temp*, *rain*, *#female*, and *male* as explanatory variables. Faecal *GCM* concentration can vary during the day [49] and between pregnant and non-pregnant females [64]. We accounted for this variation by incorporating two fixed binary variables: sample collected during morning hours (5-12:00) vs. sample collected in the afternoon (15-20:00, *collect*) and female pregnant vs. not pregnant (*preg*). As above, we treated *indID* nested in *coalID* as random intercept terms.

2a) *Emigrant body mass during dispersal stages*: In this analysis, we focused solely on emigrants, ignoring returners and residents, as we were interested in the effect of dispersal stage, which is not available for returners and residents. As above, we used $(m_t - m_0)/(m_0 * \Delta t)$ as response variable and incorporated m_0 as a fixed explanatory variable. To investigate the differences in mass loss between dispersal stages, we included a binary variable indicating post-eviction and transience stage (*stage*). Here, Δt was the time elapsed since the start of the respective dispersal stage. In contrast to the first model, we removed the variable *#pup* because no pups were present during dispersal. We further included a binary variable (*male*) indicating if at least one unrelated male had joined the coalition. For the rest, we included the same explanatory variables used in the first model: *age*, $age^{1/2}$, *temp*, *rain*, and *#female*. *IndID* was nested in *coalID* as random intercept terms.

2b) *Emigrant fGCM during dispersal stages*: As above, we only used data on emigrants. We used $\log(fGCM)$ as response variable and included the variables *stage*, *collect*, *preg*, Δt , *age*, *temp*, *rain*, and *#female*. We treated *indID* nested in *coalID* as random intercept terms.

Results

Females were evicted from their natal groups either alone ($n = 33$) or as several females at a time forming same-sex dispersing coalitions ($n = 32$) that varied in size from two to six related females (19 two-, 7 three-, 3 four-, 2 five-, and 1 six-member coalition). On the day of eviction, females were between nine months and five years

297 of age. After eviction, emigrants ($n = 26$) dispersed for an average period of 41 days
 298 (interquartile range: 13–65 days) before they settled, and returners ($n = 39$) spent an
 299 average period of 24 days (7–30 days) away from their natal group.

300 *1a) Body mass in emigrants, returners, and residents:* At eviction, emigrants were
 301 heavier ($716 \text{ g} \pm 21 \text{ SE}$) than returners ($672 \pm 26 \text{ g}$; ANOVA: Est = 43.4, SE = 8.3, p
 302 < 0.001) and resident subordinates ($585 \pm 15 \text{ g}$; ANOVA: Est = 129.9, SE = 6.8, p $<$
 303 0.001). Based on model predictions, emigrants had an average negative daily growth
 304 rate (-0.09% of body mass = $-0.61 \pm 1.33 \text{ g}$), whereas resident subordinates (0.13% ,
 305 $0.79 \pm 0.73 \text{ g}$) and returners (0.03% , $0.23 \pm 2.32 \text{ g}$) had a positive growth rate. Daily
 306 growth rates decreased with increasing initial body mass (Est = -0.003 , SE = 0.001, p
 307 = 0.002; Table 1), and the effect of initial body mass was strongest in returners (Est =
 308 -0.002 , SE = 0.001, $p = 0.006$; Figure 1A, Table 1). Young females gained body mass
 309 disproportionately faster than old females (nonlinear age effect: Est = 0.002, SE =
 310 0.001, $p = 0.012$), and females lost body mass faster at high temperatures (Est = -
 311 0.001, SE = 0.0002, $p = 0.009$). Growth rates remained constant throughout the entire
 312 dispersal event and were not influenced by coalition size.

313 *1b) fGCM in emigrants, returners, and residents:* Faecal GCM levels varied among
 314 dispersal strategies. Emigrants and returners experienced overall similar fGCM levels
 315 (Est = -0.21 , SE = 0.17, $p = 0.214$) that were on average higher than those of residents
 316 (emigrants: Est = 0.48, SE = 0.18, $p = 0.008$; returners: Est = 0.70, SE = 0.21, $p =$
 317 0.001; Figure 1B, Table 1). Faecal GCM levels increased with time after eviction (Est
 318 = 0.18, SE = 0.07, $p = 0.009$) and decreased with increasing daily temperatures (Est =
 319 -0.23 , SE = 0.07, $p < 0.001$). Faecal GCM levels did not change with changing
 320 coalition size.

321 *2a) Emigrant body mass during dispersal stages:* Mass loss of emigrants did not vary
 322 between post-eviction and transience, nor with coalition size or presence of unrelated
 323 males (Table 1). Mass at eviction had a negative effect on changes in emigrant body
 324 mass (Est = -0.003 , SE = 0.001, $p = 0.022$). Time spent in a dispersal stage did not
 325 influence emigrant body mass (Table 1).

326 *2b) Emigrant fGCM during dispersal stages:* Faecal GCM levels of emigrants varied
 327 among dispersal stages (Figure 2) but did not depend on the time emigrants spent in a

stage (Table 1). Emigrant females had higher fGCM levels during transience than during post-eviction (Est = 0.37, SE = 0.16, $p = 0.026$). Dispersing coalition size and presence of males did not affect fGCM levels (Table 1).

In all models, much of the observed variation was explained by the random effects, which is shown by the difference between the conditional and marginal R squared (Table 1). This was mostly due to high variability in daily body mass changes within each individual and between coalitions (Table 2, models 1a and 2a), and due to high variability in fGCM levels within individuals and coalitions (Table 2, models 1b and 2b). In general, variability between individuals was low (Table 2).

Discussion

Our study emphasizes that dispersal is a costly process and that costs vary between dispersal strategies and dispersal stages. Dispersing females who successfully settled lost body mass, thus confirming the expectation that loss of body mass is associated with dispersal [3]. We further showed that the negative effect of mass at eviction on daily growth rates was more pronounced in dispersers that returned to their natal group than in dispersers that successfully settled in a new territory. Finally, we showed that dispersing females experienced higher faecal glucocorticoid metabolite (fGCM) levels than their resident counterparts, especially in the later stages of dispersal. Our findings thus provide empirical support for the prediction that dispersal is associated with increased stress hormone levels [3].

The fact that dispersers that successfully settled (i.e., emigrants) were on average heavier (also reported in [41]) and less affected by the negative effect of mass at eviction than dispersers that returned to their natal group (i.e., returners), suggests that heavier individuals cope better than lighter individuals when away from the natal group. In actively dispersing species (e.g., birds and mammals), heavier individuals or individuals in better condition often have an advantage during dispersal as they can move faster and further away from the natal territory [65,66]. This increases the likelihood of finding less-saturated territories to settle and to increase breeding opportunities [2,67]. In returners, we did not observe loss of body mass during the time they spent away from the group. This is in line with a study on red squirrels

showing that individuals experience only minor changes in body mass during temporary forays around the natal site [23]. While individuals roaming through familiar areas close to the natal territory are likely to maintain their condition, permanent dispersers may suffer from inefficient foraging in unknown areas [11]. Based on the observed differences in growth rates between emigrants and returners, conclusions regarding a successful dispersal process should be drawn very carefully when based only on prospecting individuals.

The stronger negative effect of body mass at eviction on growth rates of returners may be due to their inexperience of being outside of the group. Over the first years of their life, subordinate females are generally evicted several times before they finally emigrate and establish their own group [26]. During these early evictions, young females can gain information on their surroundings (N. Maag, *pers. obs.*). Similar pre-dispersal movements are common in many species as individuals can gain information on mate availability or quality of future breeding habitat [68,69]. Pre-dispersal forays can thus reduce the costs of dispersal and improve settlement success [28,70]. We propose that lighter and less experienced individuals may not be able to survive dispersal and settle successfully, but that they collect information essential for future dispersal.

Contrary to expectations [3,24], we did not observe a decrease in growth rate with increasing dispersal time nor with time spent in a dispersal stage [3,24]. Hence, the energetic costs of being away from the group may not be as high as previously thought [3,11]. However, to assess the influence of dispersal on body condition, an investigation of distance covered by each individual could be more informative [3]. While our main interest was to investigate the variation of costs among dispersal strategies and stages, we suggest that future studies should explore the influence of dispersal distance on changes in individual body mass and stress levels. This would increase our understanding of how costs vary between different movement strategies followed by each individual.

We could not confirm previous results showing that mass loss in dispersing meerkats can be alleviated in larger coalitions [38]. However, the coalition size effect was close to being significant (Table 1, model 1a) and more information on large coalitions may have improved the model fit. Since cooperative birds and mammals can partition

cooperative behaviour like vigilance among group members [71,72], dispersing in larger coalitions is likely to reduce the individual costs of such behaviour during dispersal. This allows individuals in larger coalitions to allocate more time to foraging [38,71,72]. If increased foraging time in larger coalitions buffers against daily individual mass loss, many cooperative species that disperse in multiple-member coalitions – such as lions (*Panthera leo*), wild dogs (*Lycaon pictus*), green woodhoopoes (*Phoeniculus purpureus*), or Arabian babblers (*Turdoides squamiceps*) [13,32–34] – may experience reduced mass loss during dispersal. In addition, individuals in larger coalitions may have an advantage as they can establish a new breeding unit with a larger initial group size. Starting a new group with several helpers will likely increase total group reproduction and survival of individuals due to improved group augmentation [73,74].

Aggression from the dominant female during eviction and the lack of protection offered by the group after eviction may be responsible for the higher fGCM levels during the post-eviction stage [16]. Aggressive attacks are used by dominants of several cooperative species to render their subordinates infertile through chronic stress [75,76] and are a likely explanation for increased fGCM concentrations in meerkats [16]. Changes in physical and social environment associated with dispersal, such as unfamiliarity with the landscape and aggression from unrelated residents, are likely to intensify during transience and offer an explanation for the increased stress hormone levels during this stage of the dispersal process [77,78]. Many species cope with unpredictable stimuli by exhibiting a stress response, which may lead to increased glucocorticoid levels [6]. Increased glucocorticoid levels as response to homeostatic challenges are at first place adaptive [79]. Only when stress hormones are elevated over prolonged periods (i.e., chronically) they have deleterious impacts on an individual's fitness [80]. As such, the observed change in fGCM concentrations may be an adaptive response to the challenge of dispersal rather than a real cost of affecting fitness negatively. However, if individuals are unable to find suitable territory for settlement in time, chronic stress may lead to decreased fitness later in life [78].

In contrast to previous work by Young [38], we did not observe a reduction of stress hormone levels in females of larger coalitions. This difference is possibly due to the fact that Young [38] focused mainly on the early phases of dispersal. We showed that stress hormone levels increased over time and when moving from the post-eviction to the transience stage. Hence, the increased stress response associated with the late stages of dispersal [3] may have masked the positive effect of coalition size. In addition, studies on other species have shown that glucocorticoid levels of reproductive competitors increase during times of social instability [81,82] and such instability is likely to occur during the late stages of dispersal, when individuals establish new breeding groups. As aggression and reproductive conflict increase with group size in social species [83,84], increased reproductive competition in larger coalitions may increase individual stress hormone levels and counteract the benefits of dispersing with several helpers. Our results suggest that environmental and social circumstances can change during the dispersal process and cause variation in fGCM concentrations.

In conclusion, we show that dispersing females who successfully settle experience greater mass loss than resident females and evicted females who return to the natal group. Both emigrants and returners have higher stress hormone output than residents, especially during the later stages of dispersal. We thereby confirm previous findings, but also extend those by showing how body condition varies between different female strategies and dispersal stages. Body mass and stress hormone output seem suitable parameters for monitoring the costs associated with dispersal, as these traits often play an important role in mediating the effects of the environment on other life-history events such as survival and breeding [7–9]. A better quantification of how such costs change in response to social and environmental conditions across different stages of dispersal can therefore help gain a better mechanistic understanding of this important life history event. As many vertebrate species are expected to experience condition loss during dispersal [2,3], our results – showing how body condition changes during the later stages of dispersal – should hold implications for a wide range of taxa. Quantification of the effects of dispersal on individual condition will permit a trait-based investigation of the associated demographic rates and how these affect population dynamics.

454 **Acknowledgements**

455 Funding for this study was provided by the Swiss National Science Foundation
456 (Project CR32I3_159743). We thank the Northern Cape Conservation Authority for
457 permission to conduct this research, and the farmers neighbouring the Kuruman River
458 Reserve for granting us access to their private lands. We thank the field managers and
459 volunteers for facilitating field work and helping with data collection, in particular
460 David Gaynor and Tim Vink. Special thanks go to our field assistants David Seager,
461 Ana Morales González, Héctor Ruiz Villar, Peter Clark, Luc Le Grand, Louis Bliard,
462 Natasha Harrison, and Frances Mullany. We thank Stefanie Ganswindt for hormone
463 extraction and Andrea Heistermann for hormone analysis. We thank Luca Börger for
464 fruitful discussions and comments on the manuscript, and Tina Cornioley and Mollie
465 Brooks for their help with statistical analyses. Our research relied on records of
466 individual identities and life histories of meerkats, and research facilities maintained
467 by the Kalahari Meerkat Project, which has been supported by the European Research
468 Council (Research Grant No 294494 to T.H. Clutton-Brock), the University of Zurich
469 and the Mammal Research Institute at the University of Pretoria.

470 **Figure legends and tables**

471 **Figure 1.** Effects of (A) initial body mass at eviction on net proportional daily change
472 in body mass, and (B) dispersal time on faecal glucocorticoid metabolite (fGCM)
473 concentration in female meerkats. A) Points show average daily mass changes for
474 each female with standard errors. The lines show model predictions for each female
475 strategy (resident, returner, emigrant) when all other model predictors were set to
476 their mean. B) Points show observed fGCM concentrations and lines show model
477 predictions for each female strategy when all other model predictors were set to their
478 mean. The slopes capture the change in the response for a one-standard-deviation
479 increase in the respective variable. Significance values are given in Table 1.

480 **Figure 2.** Faecal glucocorticoid metabolite (fGCM) concentrations in female meerkat
481 emigrants depending on whether they were in the post-eviction or transience stage.
482 Empty symbols show observed fGCM concentrations and filled symbols show model
483 estimates with 95% confidence intervals.

484 **Table 1.** Effects of individual, social, and environmental factors on net proportional
 485 daily changes in body mass and faecal glucocorticoid metabolite (fGCM)
 486 concentrations of female meerkats. We report the estimate (Est), standard error (SE),
 487 significance (p, based in Wald statistics), and relative importance (RI) for each term
 488 based on the model average from models with $\Delta AICc < 2$. Marginal R^2 represents the
 489 variance explained by fixed factors (R^2_m), and conditional R^2 represents the variance
 490 explained by both fixed and random factors (R^2_c). The explanatory variables are m_0 =
 491 initial mass, $temp$ = max daily temperature, $\#female$ = average number of females,
 492 $strategy$ = emigrants vs. returners vs. residents, age = age in month, $age^{1/2}$ = sqrt of
 493 age to account for nonlinearity in growth, $rain$ = rain sum of previous 3 months, Δt =
 494 days since eviction, $\#pup$ = number of offspring < 3 months of age, $collect$ = sample
 495 collected am or pm, $preg$ = female pregnant or not, and $stage$ = post-eviction vs.
 496 transience.

Model	Est	SE	p	RI
1a) Body mass in emigrants, returners, and residents			$R^2_m=0.13$, $R^2_c=0.72$	
<i>intercept</i>	0.001	0.001	0.371	
<i>m₀</i>	-0.003	0.001	0.002	1.00
<i>temp</i>	-0.001	0.000	0.009	1.00
<i>#female</i>	-0.002	0.001	0.085	1.00
<i>strategyReturner</i>	0.000	0.002	0.842	0.70
<i>strategyEmigrant</i>	0.000	0.002	0.857	
<i>strategyReturner:m₀</i>	-0.003	0.001	0.006	0.70
<i>strategyEmigrant:m₀</i>	-0.001	0.001	0.339	
<i>age^{1/2}</i>	0.002	0.001	0.012	0.53
<i>age</i>	0.002	0.001	0.013	0.47
<i>rain</i>	0.001	0.001	0.201	0.42
<i>Δt</i>	0.000	0.000	0.243	0.31
<i>strategyReturner:#female</i>	-0.001	0.004	0.909	0.10
<i>strategyEmigrant:#female</i>	-0.004	0.002	0.104	
<i>#pup</i>	0.000	0.000	0.635	0.04
1b) fGCM in emigrants, returners, and residents			$R^2_m=0.15$, $R^2_c=0.29$	
<i>intercept</i>	5.180	0.150	<0.001	
<i>collectPM</i>	-0.405	0.111	<0.001	fixed
<i>pregPregnant</i>	0.138	0.129	0.284	fixed
<i>temp</i>	-0.231	0.065	<0.001	1.00
<i>strategyReturner</i>	0.697	0.205	0.001	1.00
<i>strategyEmigrant</i>	0.482	0.180	0.008	
<i>Δt</i>	0.180	0.068	0.009	1.00
<i>#female</i>	0.150	0.081	0.064	0.75
<i>age</i>	0.087	0.080	0.282	0.28
2a) Emigrant body mass during dispersal stages			$R^2_m=0.09$, $R^2_c=0.70$	
<i>intercept</i>	0.001	0.002	0.760	
<i>m₀</i>	-0.003	0.001	0.022	0.88
<i>stageTransience</i>	-0.001	0.001	0.293	0.18
<i>temp</i>	0.000	0.001	0.389	0.15
<i>rain</i>	0.001	0.001	0.550	0.12
<i>#female</i>	-0.001	0.002	0.544	0.12
2b) Emigrant fGCM during dispersal stages			$R^2_m=0.11$, $R^2_c=0.25$	
<i>intercept</i>	5.611	0.119	<0.001	
<i>collectPM</i>	-0.447	0.131	0.001	fixed
<i>pregPregnant</i>	0.054	0.148	0.717	fixed
<i>stageTransience</i>	0.366	0.163	0.026	1.00
<i>rain</i>	-0.106	0.073	0.148	0.44
<i>temp</i>	-0.094	0.071	0.189	0.41
<i>age</i>	0.058	0.083	0.488	0.11

Table 2. Within-individual and within-coalition repeatability (R) for each model. R < 0.40 means low variability between individuals (coalitions, respectively,) and high variability within individuals (coalitions, respectively). R > 0.60 means high variability between individuals (coalitions, respectively) and low variability within individuals (coalitions, respectively) [63].

Model	Random factor	R
1a) Body mass in emigrants, returners, and residents	Individual	0.33
	Coalition	0.62
1b) fGCM in emigrants, returners, and residents	Individual	0.02
	Coalition	0.15
2a) Emigrant body mass during dispersal stages	Individual	0.10
	Coalition	0.65
2b) Emigrant fGCM during dispersal stages	Individual	0.00
	Coalition	0.16

505 **References**

- 506 1. Bowler DE, Benton TG. 2005 Causes and consequences of animal dispersal strategies:
507 relating individual behaviour to spatial dynamics. *Biol. Rev. Camb. Philos. Soc.* **80**, 205–
508 225.
- 509 2. Clobert J, Baguette M, Benton TG, Bullock JM. 2012 *Dispersal Ecology and Evolution*.
510 Oxford University Press.
- 511 3. Bonte D *et al.* 2012 Costs of dispersal. *Biol. Rev. Camb. Philos. Soc.* **87**, 290–312.
- 512 4. Nevoux M, Arlt D, Nicoll M, Jones C, Norris K. 2013 The short-and long-term fitness
513 consequences of natal dispersal in a wild bird population. *Ecol. Lett.* **16**, 438–445.
- 514 5. Kingma SA, Komdeur J, Burke T, Richardson DS. 2017 Differential dispersal costs and
515 sex-biased dispersal distance in a cooperatively breeding bird. *Behav. Ecol.* **28**, 1113–
516 1121.
- 517 6. Romero LM. 2004 Physiological stress in ecology: lessons from biomedical research.
518 *Trends Ecol. Evol.* **19**, 249–255.
- 519 7. Crespi EJ, Williams TD, Jessop TS, Delehanty B. 2013 Life history and the ecology of
520 stress: how do glucocorticoid hormones influence life-history variation in animals?
521 *Funct. Ecol.* **27**, 93–106.
- 522 8. Plard F, Gaillard J-M, Coulson T, Delorme D, Warnant C, Michallet J, Tuljapurkar S,
523 Krishnakumar S, Bonenfant C. 2015 Quantifying the influence of measured and
524 unmeasured individual differences on demography. *J. Anim. Ecol.* **84**, 1434–1445.
- 525 9. Ozgul A, Childs DZ, Oli MK, Armitage KB, Blumstein DT, Olson LE, Tuljapurkar S,
526 Coulson T. 2010 Coupled dynamics of body mass and population growth in response to
527 environmental change. *Nature* **466**, 482–485.
- 528 10. Pride ER. 2005 High faecal glucocorticoid levels predict mortality in ring-tailed lemurs
529 (*Lemur catta*). *Biol. Lett.* **1**, 60–63.
- 530 11. Pinter-Wollman N, Isbell LA, Hart LA. 2009 The relationship between social behaviour
531 and habitat familiarity in African elephants (*Loxodonta africana*). *Proc. Biol. Sci.* **276**,
532 1009–1014.
- 533 12. Metzgab LH. 1967 An Experimental Comparison of Screech Owl Predation on Resident
534 and Transient White-Footed Mice (*Peromyscus Leucopus*). *J. Mammal.* **48**, 387–391.
- 535 13. Packer C, Pusey AE. 1982 Cooperation and competition within coalitions of male lions:
536 kin selection or game theory? *Nature* **296**, 740.
- 537 14. Boydston EE, Morelli TL, Holekamp KE. 2001 Sex Differences in Territorial Behavior
538 Exhibited by the Spotted Hyena (*Hyaenidae, Crocuta crocuta*). *Ethology* **107**, 369–385.
- 539 15. Solomon NG. 2003 A reexamination of factors influencing philopatry in rodents. *J.*
540 *Mammal.* **84**, 1182–1197.
- 541 16. Young AJ, Carlson AA, Monfort SL, Russell AF, Bennett NC, Clutton-Brock T. 2006
542 Stress and the suppression of subordinate reproduction in cooperatively breeding
543 meerkats. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 12005–12010.


- 544 17. Soulsbury CD, Baker PJ, Iossa G, Harris S. 2008 Fitness costs of dispersal in red foxes
545 (*Vulpes vulpes*). *Behav. Ecol. Sociobiol.* **62**, 1289–1298.
- 546 18. Calleri DV, Rosengaus RB, Traniello JFA. 2006 Disease and colony establishment in the
547 dampwood termite *Zootermopsis angusticollis*: survival and fitness consequences of
548 infection in primary reproductives. *Insectes Soc.* **53**, 204–211.
- 549 19. Adamo SA, Roberts JL, Easy RH, Ross NW. 2008 Competition between immune
550 function and lipid transport for the protein apolipoprotein III leads to stress-induced
551 immunosuppression in crickets. *J. Exp. Biol.* **211**, 531–538.
- 552 20. Srygley RB, Lorch PD, Simpson SJ, Sword GA. 2009 Immediate protein dietary effects
553 on movement and the generalised immunocompetence of migrating Mormon crickets
554 *Anabrus simplex* (Orthoptera: Tettigoniidae). *Ecol. Entomol.* **34**, 663–668.
- 555 21. Cant MA, Otali E, Mwanguhya F. 2001 Eviction and dispersal in co-operatively
556 breeding banded mongooses (*Mungos mungo*). *J. Zool.* **254**, 155–162.
- 557 22. Vanderwerf EA. 2008 Sources of Variation in Survival, Recruitment, and Natal
558 Dispersal of the Hawai'i 'Elepaio. *Condor* **110**, 241–250.
- 559 23. Larsen KW, Boutin S. 1994 Movements, Survival, and Settlement of Red Squirrel
560 (*Tamiasciurus Hudsonicus*) Offspring. *Ecology* **75**, 214–223.
- 561 24. Ridley AR, Raihani NJ, Nelson-Flower MJ. 2008 The cost of being alone: the fate of
562 floaters in a population of cooperatively breeding pied babblers *Turdoides bicolor*. *J.*
563 *Avian Biol.* **39**, 389–392.
- 564 25. Young AJ, Carlson AA, Clutton-Brock T. 2005 Trade-offs between extraterritorial
565 prospecting and helping in a cooperative mammal. *Anim. Behav.* **70**, 829–837.
- 566 26. Clutton-Brock TH, Brotherton PN, Smith R, McIlrath GM, Kansky R, Gaynor D,
567 O'Riain MJ, Skinner JD. 1998 Infanticide and expulsion of females in a cooperative
568 mammal. *Proc. Biol. Sci.* **265**, 2291–2295.
- 569 27. Young AJ, Monfort SL. 2009 Stress and the costs of extra-territorial movement in a
570 social carnivore. *Biol. Lett.* **5**, 439–441.
- 571 28. Haughland DL, Larsen KW. 2004 Exploration correlates with settlement: red squirrel
572 dispersal in contrasting habitats. *J. Anim. Ecol.* **73**, 1024–1034.
- 573 29. Brown JL, Brown ER, Brown SD, Dow DD. 1982 Helpers: effects of experimental
574 removal on reproductive success. *Science* **215**, 421–422.
- 575 30. Courchamp F, Grenfell BT, Clutton-Brock TH. 2000 Impact of natural enemies on
576 obligately cooperative breeders. *Oikos* **91**, 311–322.
- 577 31. Clutton-Brock TH, Russell AF, Sharpe LL, Brotherton PN, McIlrath GM, White S,
578 Cameron EZ. 2001 Effects of helpers on juvenile development and survival in meerkats.
579 *Science* **293**, 2446–2449.
- 580 32. Ligon JD, Ligon SH. 1978 Communal breeding in green woodhoopoes as a case for
581 reciprocity. *Nature* **276**, 496.
- 582 33. Lundy KJ, Parker PG, Zahavi A. 1998 Reproduction by subordinates in cooperatively
583 breeding Arabian babblers is uncommon but predictable. *Behav. Ecol. Sociobiol.* **43**,
584 173–180.

- 585 34. McNutt JW. 1996 Sex-biased dispersal in African wild dogs, *Lycaon pictus*. *Anim.*
586 *Behav.* **52**, 1067–1077.
- 587 35. Clutton-Brock TH, Gaynor D, McIlrath GM. 1999 Predation, group size and mortality in
588 a cooperative mongoose, *Suricata suricatta*. *Journal of Animal*
- 589 36. Packer C, Scheel D, Pusey AE. 1990 Why Lions Form Groups: Food is Not Enough.
590 *Am. Nat.* **136**, 1–19.
- 591 37. Wilson ML, Britton NF, Franks NR. 2002 Chimpanzees and the mathematics of battle.
592 *Proceedings of the Royal Society B: Biological Sciences* **269**, 1107–1112.
- 593 38. Young AJ. 2004 Subordinate tactics in cooperative meerkats: helping, breeding and
594 dispersal. University of Cambridge.
- 595 39. Griffin AS, Pemberton JM, Brotherton PNM, McIlrath G, Gaynor D, Kansky R, O’Riain
596 J, Clutton-Brock TH. 2003 A genetic analysis of breeding success in the cooperative
597 meerkat (*Suricata suricatta*). *Behav. Ecol.* **14**, 472–480.
- 598 40. Maag N, Cozzi G, Clutton-Brock T, Ozgul A. 2018 Density-dependent dispersal
599 strategies in a cooperative breeder. *Ecology* (doi:10.1002/ecy.2433)
- 600 41. Ozgul A, Bateman AW, English S, Coulson T, Clutton-Brock TH. 2014 Linking body
601 mass and group dynamics in an obligate cooperative breeder. *J. Anim. Ecol.* **83**, 1357–
602 1366.
- 603 42. Jordan NR, Cherry MI, Manser MB. 2007 Latrine distribution and patterns of use by
604 wild meerkats: implications for territory and mate defence. *Anim. Behav.* **73**, 613–622.
- 605 43. Cozzi G, Maag N, Börger L, Clutton-Brock TH, Ozgul A. 2018 Socially informed
606 dispersal in a territorial cooperative breeder. *J. Anim. Ecol.* **87**, 838–849.
- 607 44. Börger L, Fryxell JM. 2012 Quantifying individual differences in dispersal using net
608 squared displacement. In *Dispersal Ecology and Evolution* (eds J Clobert, M Baguette,
609 TG Benton, JM Bullock), pp. 222–230. Oxford University Press, Oxford, UK.
- 610 45. Cozzi G, Chynoweth M, Kusak J, Çoban E, Çoban A, Ozgul A, Şekercioğlu ÇH. 2016
611 Anthropogenic food resources foster the coexistence of distinct life history strategies:
612 year-round sedentary and migratory brown bears. *J. Zool.* **300**, 142–150.
- 613 46. Clutton-Brock TH *et al.* 1998 Costs of cooperative behaviour in suricates (*Suricata*
614 *suricatta*). *Proc. Biol. Sci.* **265**, 185–190.
- 615 47. Doolan’ SP, Macdonald DW. 1997 Breeding and juvenile survival among slender-tailed
616 meerkats (*Suricatu suricatta*) in the south-western Kalahari: ecological and social
617 influences. *J. Zool., Lond.* **242**, 309–327.
- 618 48. Sharp SP, English S, Clutton-Brock TH. 2013 Maternal investment during pregnancy in
619 wild meerkats. *Evol. Ecol.* **27**, 1033–1044.
- 620 49. Braga Goncalves I, Heistermann M, Santema P, Dantzer B, Mausbach J, Ganswindt A,
621 Manser MB. 2016 Validation of a Fecal Glucocorticoid Assay to Assess Adrenocortical
622 Activity in Meerkats Using Physiological and Biological Stimuli. *PLoS One* **11**,
623 e0153161.
- 624 50. Heistermann M, Finke M, Hodges JK. 1995 Assessment of female reproductive status in
625 captive-housed Hanuman langurs (*Presbytis entellus*) by measurement of urinary and

- fecal steroid excretion patterns. *Am. J. Primatol.* **37**, 275–284.
51. Ganswindt A, Palme R, Heistermann M, Borragan S, Hodges JK. 2003 Non-invasive assessment of adrenocortical function in the male African elephant (*Loxodonta africana*) and its relation to musth. *Gen. Comp. Endocrinol.* **134**, 156–166.
52. Mausbach J, Goncalves IB, Heistermann M, Ganswindt A, Manser MB. 2017 Meerkat close calling patterns are linked to sex, social category, season and wind, but not fecal glucocorticoid metabolite concentrations. *PLoS One* **12**, e0175371.
53. Heistermann M, Ademmer C, Kaumanns W. 2004 Ovarian Cycle and Effect of Social Changes on Adrenal and Ovarian Function in *Pygathrix nemaeus*. *Int. J. Primatol.* **25**, 689–708.
54. Bates D, Maechler M, Bolker B, Walker S, Others. 2014 lme4: Linear mixed-effects models using Eigen and S4. *R package version 1*, 1–23.
55. R Core Team. 2013 R: A language and environment for statistical computing.
56. Bartoń K. 2018 MuMIn: multi-model inference. R package version 1.40.4. <http://CRAN.R-project.org/package=MuMIn>
57. Akaike H. 1998 Information Theory and an Extension of the Maximum Likelihood Principle. In *Selected Papers of Hirotugu Akaike* (eds E Parzen, K Tanabe, G Kitagawa), pp. 199–213. New York, NY: Springer New York.
58. Burnham, K.P. & Anderson, D.R. 2002. Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach. Berlin, Germany: Springer.
59. Belsley DA, Kuh E, Welsch RE. 2005 *Regression Diagnostics: Identifying Influential Data and Sources of Collinearity*. John Wiley & Sons.
60. Pinheiro JC, Bates DM, editors. 2000 Linear Mixed-Effects Models: Basic Concepts and Examples. In *Mixed-Effects Models in S and S-PLUS*, pp. 3–56. New York, NY: Springer New York.
61. Calder WA. 1984 *Size, Function, and Life History*. Courier Corporation.
62. English S, Bateman AW, Clutton-Brock TH. 2012 Lifetime growth in wild meerkats: incorporating life history and environmental factors into a standard growth model. *Oecologia* **169**, 143–153.
63. Stoffel MA, Nakagawa S, Schielzeth H. 2017 rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods Ecol. Evol.* **8**, 1639–1644.
64. Sheriff MJ, Dantzer B, Delehanty B, Palme R, Boonstra R. 2011 Measuring stress in wildlife: techniques for quantifying glucocorticoids. *Oecologia* **166**, 869–887.
65. Delgado M del M, Penteriani V, Revilla E, Nams VO. 2010 The effect of phenotypic traits and external cues on natal dispersal movements. *J. Anim. Ecol.* **79**, 620–632.
66. Debeffe L, Morellet N, Cargnelutti B, Lourtet B, Bon R, Gaillard J-M, Mark Hewison AJ. 2012 Condition-dependent natal dispersal in a large herbivore: heavier animals show a greater propensity to disperse and travel further. *J. Anim. Ecol.* **81**, 1327–1327.
67. Gundersen G, Andreassen HP, Ims RA. 2002 Individual and population level

- 666 determinants of immigration success on local habitat patches: an experimental approach.
667 *Ecol. Lett.* **5**, 294–301.
- 668 68. Davidian E, Courtiol A, Wachter B, Hofer H, Höner OP. 2016 Why do some males
669 choose to breed at home when most other males disperse? *Sci Adv* **2**, e1501236.
- 670 69. Clobert J, Le Galliard J-F, Cote J, Meylan S, Massot M. 2009 Informed dispersal,
671 heterogeneity in animal dispersal syndromes and the dynamics of spatially structured
672 populations. *Ecol. Lett.* **12**, 197–209.
- 673 70. Stamps JA, Krishnan VV. 1999 A Learning-Based Model of Territory Establishment. *Q.*
674 *Rev. Biol.* **74**, 291–318.
- 675 71. Gaston AJ. 1977 Social behaviour within groups of jungle babblers (*Turdoides striatus*).
676 *Anim. Behav.* **25**, 828–848.
- 677 72. Elgar MA. 1989 Predator vigilance and group size in mammals and birds: a critical
678 review of the empirical evidence. *Biol. Rev. Camb. Philos. Soc.* **64**, 13–33.
- 679 73. Kokko H, Johnstone RA, Clutton-Brock TH. 2001 The evolution of cooperative
680 breeding through group augmentation. *Proc. Biol. Sci.* **268**, 187–196.
- 681 74. Clutton-Brock T. 2002 Breeding together: kin selection and mutualism in cooperative
682 vertebrates. *Science* **296**, 69–72.
- 683 75. Wingfield JC, Hegner RE, Lewis DM. 1991 Circulating levels of luteinizing hormone
684 and steroid hormones in relation to social status in the cooperatively breeding white-
685 browed sparrow weaver, *Plocepasser mahali*. *J. Zool.* **225**, 43–58.
- 686 76. Schoech SJ, Mumme RL, Moore MC. 1991 Reproductive Endocrinology and
687 Mechanisms of Breeding Inhibition in Cooperatively Breeding Florida Scrub Jays
688 (*Aphelocoma c. coerulescens*). *Condor* **93**, 354–364.
- 689 77. Rubenstein DR. 2007 Stress hormones and sociality: integrating social and
690 environmental stressors. *Proc. Biol. Sci.* **274**, 967–975.
- 691 78. Creel S, Dantzer B, Goymann W, Rubenstein DR. 2013 The ecology of stress: effects of
692 the social environment. *Funct. Ecol.* **27**, 66–80.
- 693 79. McEwen BS, Wingfield JC. 2003 The concept of allostasis in biology and biomedicine.
694 *Horm. Behav.* **43**, 2–15.
- 695 80. Wingfield JC, Maney DL, Breuner CW, Jacobs JD, Lynn S, Ramenofsky M, Richardson
696 R. 1998 Ecological Bases of Hormone—Behavior Interactions: The ‘Emergency Life
697 History Stage’. *Integr. Comp. Biol.* **38**, 191–206.
- 698 81. Sapolsky RM. 1993 The physiology of dominance in stable versus unstable social
699 hierarchies. In *Primate social conflict* (eds WA Mason, SP Mendoza), pp. 171–204.
700 State University of New York Press, Albany, New York, USA.
- 701 82. Creel S. 2001 Social dominance and stress hormones. *Trends Ecol. Evol.* **16**, 491–497.
- 702 83. Dantzer B, Bennett NC, Clutton-Brock TH. 2017 Social conflict and costs of
703 cooperation in meerkats are reflected in measures of stress hormones. *Behav. Ecol.* **28**,
704 1131–1141.
- 705 84. Clutton-Brock TH, Hodge SJ, Flower TP, Spong GF, Young AJ. 2010 Adaptive

706 suppression of subordinate reproduction in cooperative mammals. *Am. Nat.* **176**, 664–
707 673.



Log fGCM concentration

8
7
6
5
4
3

Post-Eviction

Transience

Dispersal stage

 $p = 0.026$ <http://mc.manuscriptcentral.com/prsb>